

Evidence-Based Treatment for Melasma: Expert Opinion and a Review

Krupa Shankar · Kiran Godse · Sanjeev Aurangabadkar · Koushik Lahiri · Venkat Mysore ·
Anil Ganjoo · Maya Vedomurty · Malavika Kohli · Jaishree Sharad · Ganesh Kadhe ·
Pashmina Ahirrao · Varsha Narayanan · Salman Abdulrehman Motlekar

To view enhanced content go to www.dermtherapy-open.com

Received: July 7, 2014 / Published online: October 1, 2014

© The Author(s) 2014. This article is published with open access at Springerlink.com

ABSTRACT

Introduction: Melasma is one of the most common pigmentary disorders seen by dermatologists and often occurs among women with darker complexion (Fitzpatrick skin type IV–VI). Even though melasma is a

widely recognized cause of significant cosmetic disfigurement worldwide and in India, there is a lack of systematic and clinically usable treatment algorithms and guidelines for melasma management. The present article outlines the epidemiology of melasma, reviews the various treatment options along with their mode of action, underscores the diagnostic dilemmas and quantification of illness, and weighs the evidence of currently available therapies.

Electronic supplementary material The online version of this article (doi:[10.1007/s13555-014-0064-z](https://doi.org/10.1007/s13555-014-0064-z)) contains supplementary material, which is available to authorized users.

K. Shankar (✉)
Department of Dermatology, Skin Diagnosis Centre,
Mahabodhi Mallige Hospital, Manipal Hospital, 98,
HAL Airport Road, Bangalore 560017, Karnataka,
India
e-mail: dermakrupa@yahoo.com

K. Godse
Shree Skin Center and Pathology Lab, Nerul,
Navi Mumbai, India

S. Aurangabadkar
Skin and Laser Clinic, 1st Floor, Brij Tarang,
Green Lands, Begumpet, Hyderabad,
Andhra Pradesh, India

K. Lahiri
Mani Square, IT-7A, 7th Floor 164/1 Manickatala
Road, Kolkata, India

V. Mysore
Venkat Charnalaya Clinic, 3437, 1st G Cross,
7th Main, Subbanna Garden, Vijayanagar,
Bangalore, Karnataka, India

A. Ganjoo
Skin and Cosmetology Centre, 105/4 LSC Gujrawala
Town, Delhi (North), New Delhi 110009, India

M. Vedomurty
RSV Skin and Laser Centre, 9/5, Mahalingam,
2nd Cross Street, Mahalingapuram, Chennai,
Tamil Nadu, India

M. Kohli
Skin Secrets, 306/403/408 Doctor House, 14 Peddar
Road, Mumbai 400 026, Maharashtra, India

J. Sharad
Skinfinity Skin and Laser Clinic, Mira Belle, Above
Scandal Shoe Shop, Near National College, Linking
Road, Bandra (W), Mumbai, Maharashtra, India

G. Kadhe · P. Ahirrao · V. Narayanan ·
S. A. Motlekar
Department of Medical Affairs, Wockhardt Ltd.,
Wockhardt Towers, Bandra Kurla Complex,
Mumbai 400051, Maharashtra, India

Methods: A panel of eminent dermatologists was created and their expert opinion was sought to address lacunae in information to arrive at a working algorithm for optimizing outcome in Indian patients. A thorough literature search from recognized medical databases preceded the panel discussions. The discussions and

melasma management, including clinical assessment, investigations, drug therapies, combinations, procedural therapies, and newer emerging therapies, from peer reviewed journals were included. Articles not adequately powered or not randomized with poorly defined methodology and articles with conflicting or non-committal results were excluded.

A total of 115 publications until August 2013 were obtained by electronic database searches of which 52 were randomized controlled trials (1 retrospective) and 42 were review articles (17 were reviews and discussions of specific drug/combination or procedural treatment). The remaining publications included eight articles on prevalence data, five articles of case series, five articles on investigations and histopathology, and three articles on validating scoring and grading systems. The literature search was completed and then followed by three panel discussion sessions, each in the form of a 1-day residential focused program where clinical experiences of each panelist were brought to the table along with the study of the associated literature and publications. All discussions, suggestions, and panel consensus were recorded by a medical writer who was present. In addition, experienced persons from the field of pharmaceuticals also contributed with respect to the pharmacological aspects of the various drug therapies. In between each of these sessions, the outcome and consensus of the previous session were drafted and shared with the panel members for further refining. After all panel discussion sessions were concluded, the final draft of the panel consensus and algorithm was prepared along with the summary of the reviewed literature. This draft was finally refined and accepted by the panelists to come up with a clinically practical, relevant, and acceptable treatment algorithm for melasma. The deployment of this algorithm is expected to act

as a basis for guiding and refining therapy in the future.

This article is based on previously conducted studies and the expert opinion of the authors, and does not involve any new studies of human or animal subjects performed by any of the authors.

RISK AND PREDISPOSING FACTORS

Melasma commonly occurs in women during their reproductive years. Pregnancy and use of oral contraceptives through estrogen mediated melanocyte stimulation is apparent in the pathogenesis of melasma [2–4]. Dark-haired persons (Fitzpatrick skin types III–VI) are more susceptible to melasma [5]. Several studies have reported a high incidence of melasma in family members suggesting its genetic predisposition [3, 6–11].

The ultraviolet (UV) component of sunlight is the major triggering and aggravating factor in melasma causing focal melanocyte hyperplasia and increase in melanosomes [1, 12, 13].

The use of topical cosmetics and fragrances as well as phototoxic drugs can also trigger or aggravate melasma by sensitizing the skin and inducing postinflammatory hyperpigmentation (PIH). Melasma is considered as photocontact dermatitis in a study conducted by Verallo-Rowell [14].

Melasma is classified according to the depth of melanin pigment into epidermal melasma with mixed melasma (a combination of the epidermal and dermal types) having a worse prognosis [1].

PRINCIPLES OF THERAPY

The objectives of melasma therapy should be protection from sunlight and

CLINICAL ASSESSMENT OF MELASMA

Traditionally, Wood's lamp examination is done to identify the location of pigment, but is limited to epidermal melasma and cannot be used reliably in Fitzpatrick skin types V and VI as dark-skin melanin pigmentation obscures the detection of dermal melanin [19]. Melasma is reliably graded on the basis of area and severity parameters [i.e., melasma area and severity index (MASI) score] [20, 21].

PHARMACOLOGIC THERAPY

Sunscreens

Sunscreen use is vital throughout treatment and post-treatment to prevent hyperpigmentation and relapse of melasma. Broad-spectrum anti-UV A and B sunscreens with physical blocking agents like ZnO and TiO₂ (SPF >30) should be used regularly to cover the affected areas. Patients with Fitzpatrick IV to VI skin types [22] are at a heightened risk from sun exposure [23, 24]. Indian patients avoid sunscreens due to oily, sticky nature and heat from exothermic reaction of chemicals. However, physical sunscreens do not release heat.

Hydroquinone

Hydroquinone (HQ; dihydroxybenzene) is structurally similar to precursors of melanin. HQ inhibits the conversion of dopa to melanin by blocking tyrosinase action [25] and inhibits the formation, melanization, and degradation of melanosomes. HQ also affects the membranous structures of melanocytes and eventually causes necrosis of whole melanocytes [26].

HQ has been used to treat hyperpigmentation for more than 50 years. While controversy exists regarding the long-term safety of HQ, its efficacy in treating melasma, both alone and in combination with other agents is well established [5]. HQ 2–4% is prominently used as a mono-therapy or in a combination cream. HQ preparations >5% are not recommended due to irritation, except for refractory cases [27, 28]. Pigment lightening by HQ becomes evident after 5–7 weeks of the treatment. Treatment with HQ should be continued for at least 3 months and up to 1 year [29].

For patients on HQ therapy, regular medical follow-up is essential: every 3 months for high phototype skin (Fitzpatrick type V and VI) patients and every 6 months for lighter skin types. Common adverse events (AEs) are erythema and burning. Other rare AEs are ochronosis and confetti-like depigmentation. Speckling or reticulation indicates ochronosis. Histopathology shows short, stout, curvilinear, ochre-colored fibers in the papillary dermis [30].

Tretinoin

Tretinoin, a retinoid (RA), inhibits transcription of the key melanin synthesis enzyme tyrosinase [16–18]. Tretinoin plays an important role in the triple-combination cream (described later) and is used as a chemical peel.

Corticosteroids

Topical corticosteroids are anti-inflammatory molecules that exert an anti-metabolic effect on melanocytes, resulting in a decreased epidermal turnover and thus, may produce a mild pigment-reducing effect [31]. Corticosteroids are an active component of the triple-combination creams. Triple formulations

using different corticosteroids have shown efficacy, for example, dexamethasone [32], hydrocortisone 1% [33], mometasone [34, 35], and fluorinated steroids [22, 36–41]. Researchers found that fluorinated steroids were more effective and safer than non-fluorinated steroids, for example, 0.01% fluocinolone acetonide and fluticasone [22].

Kojic Acid

Kojic acid can substitute for HQ if a patient is intolerant to HQ. Kojic acid inhibits tyrosinase by chelating copper at the enzyme's active site. It is available in 1–4% concentrations, and also in combination with HQ. Caution is required in its use as kojic acid is a known sensitizer [5].

Azelaic Acid

Azelaic acid (AA) is anti-proliferative and selectively cytotoxic towards hyperactive melanocytes, inhibiting tyrosinase and mitochondrial oxidoreductase enzymes with minimal effects on normally pigmented skin [42]. In different trials, AA treatment was found to be less or as equally efficacious as HQ [42, 43] and hence may be used in case of intolerance to HQ.

Triple Combination

Based on the current evidence: sun avoidance, sunscreen use, and triple combination regimen is the most effective first-line treatment for melasma. Combination creams are more efficient bleaching agents than monotherapies. The combination of HQ, a RA, and a topical steroid (5% HQ, 0.1% RA, and 0.1% dexamethasone), or Kligman and Willis' formula was developed to enhance the efficacy of each individual ingredient, shortening the

duration of therapy and reducing the risk of AEs [44, 45]. Since then, some variation of this formula is the most commonly used therapy for melasma worldwide [32]. Tretinoin prevents the oxidation of HQ and improves epidermal penetration while the steroid reduces irritation from the other two ingredients and suppresses biosynthetic and secretory functions of melanocytes, leading to an early response in melasma. The synergistic action of the three topical agents achieves significantly higher depigmentation than either agent alone. Improvement is seen in 8 weeks without any significant AE [46].

In India, a cream-based combination of HQ, tretinoin and mometasone has been extensively used after its launch in 2004. Long-term use on the face (generally >12 weeks) of corticosteroids, particularly mid-potent ones like mometasone can cause skin atrophy, telangiectasias, and/or an acneiform eruption [37]. Moreover, a relapse is common when mometasone use is stopped [47]. The combination treatment is supposed to be used for a maximum of 4–8 weeks after which the treatment should be stopped or the dose gradually reduced, and finally replaced with safer treatment modalities [48]. We found mometasone-based combination with glycolic acid (GA) peels caused AEs like hyperpigmentation, irritation, and persistent erythema in 2 out of 10 patients (20%) [34]. Triple-combination creams with mometasone should be prescribed only after thorough patient counseling with explanation of long-term AEs.

Evidence-based studies have shown triple combination of 4% HQ with 0.05% tretinoin and 0.01% fluocinolone acetonide to be the most efficacious and safe available topical modality for the first-line treatment of melasma [40, 41]. It is the only ointment

currently approved by the US Food and Drug Administration for the treatment of melasma. The safety and efficacy of the combination was initially demonstrated in 641 melasma patients in a multicenter, investigator blinded, randomized prospective trial [36, 39–41]. Night use of the triple-combination cream as compared to dual-combination creams (HQ + tretinoin, HQ + fluocinolone, or tretinoin + fluocinolone) achieved complete or near-complete clearance in 77% of patients. Clinically significant improvement was noted within 4–8 weeks [39]. The most common AEs were mild local irritation, erythema, and skin peeling. For the Indian skin, which is mostly of Fitzpatrick type IV–VI, a reduced HQ combination of 2% HQ + 0.05% tretinoin + 0.01% fluocinolone acetonide (HQ + RA + FA) was launched in 2010 [47]. Safety studies in Indian population need to be established. A recent Cochrane review of 20 studies with a total of 2,125 participants with 23 different treatments concluded that triple-combination cream was significantly more effective than HQ alone (relative risk 1.58, 95% CI 1.26–1.97) or dual combination [38].

The fluocinolone-based triple cream is safe in the treatment of moderate to severe melasma for up to 24 weeks. The risk of skin atrophy after 6 months of treatment with fluocinolone acetonide 0.01%, HQ 4%, and tretinoin 0.05% cream is very low as shown by histological examination [4].

In the expert opinion of the authors, fluticasone seems more effective and safer than mometasone and fluocinolone in the triple combination due to less long-term (> 12 weeks) steroidal AEs. Once-daily application of fluticasone (0.05%) was as efficacious as twice-daily application of betamethasone (0.12%), with less skin atrophy and an absence of systemic AE of

hypothalamic–pituitary–adrenal axis suppression [37]. In another study, fluticasone did not cause cortisol suppression or major skin atrophy, and was similar to mometasone furoate in efficacy [49].

Dual Combination

The dual combinations are recommended if triple combination is not available or if patients are intolerant to it. Those available in India include HQ and GA, HQ and Kojic acid, and mequinol and tretinoin. Table 2 gives an overview of studies that provide evidence on topical agents for treating melasma.

Adjunctive Therapies

These include a wide range of chemicals and natural extracts have been tested against melasma (Table 3). Tranexamic acid is the most common adjunctive therapy to be used and works by decreasing melanogenesis in epidermal melanocytes and provides a rapid and sustained lightening in melasma [50–53]. Other therapies include a substance called ‘antipollon’, a finely grained aluminum silicate that possesses the ability to adsorb melanin. Laboratory studies with different concentrations have shown up to 86% melanin adsorption with 0.8% antipollon. However, clinical studies establishing its efficacy are awaited (Antipollon-HT efficacy testing Nikkol Chemicals, data on file).

Interventions

Intervention procedures are used in combination with topical first-line therapy to treat recalcitrant melasma when patient shows poor or no response. Intervention techniques include chemical peels, lasers, intense pulsed

T		2 E c c		P r a		P ca r a		P	
R		Site	T	i	i	P i i ,	S v i	T i i	R i i
M r a .	[77]	R, DB	4% HQ	. P ac b	+ SPF 30	48	N/K	20	38% HQ/8.3% P ac b , a P r
Ha c a c [27]		R, DB, SF	4% HQ	. P ac b	+ SPF 25	30	N/K	3	79% HQ/67% SWC P r
Far [78]		R, O	4% HQ	. a a c ac c	20%	29	N/K	2	A r 2 P r a , MASI c r a 6.2 ± 3.6 P r HQ a c 3.8 ± 2.8 P r a a c ac c
E P a -P r a . [79]		R, O, SF	4% HQ	. 5% a c r b c ac c		16	N/K	16	HQ c P r 93% c a c c r u , c P ar c P r 62.5% P r a c r b c ac c c . S c c P r P r 68.7% P r HQ . 6.2% P r a c r b c ac c
Na c a a . [80]		R, O	Pr P r 2% HQ , 0.025% RA c	ca (P r) 2 P r b r a r r P r ac c ac c P r (r 2 P r r 12 P r)		50	N/K	6	HQ u P r r RA a a P r a a a P r r u ac c P r a c c c a P r c c P r P r ac P r P r a
Ba a a c G r P [43]		R, DB, MC	4% HQ	. 20% a a c ac c		329	N/K	24	65%/73% c r c a a c/HQ P a
P q u r Mar a . [81], V r a - R a . [82]		R, DB O	4% HQ	. 20% a a c ac c		60	N/K	24	A a c ac c P a b r P a P r HQ P r r a a a
V r a -R a . [82], L [83]		R, DB	2% HQ	. 20% a a c ac c		155	N/K	24	73% P r a a c ac c P a , c P ar c P r 19% P r HQ P a , P ac c c r a r u
L [83], F r r a C a r a . [84]		R	2% KA, 2% HQ, 10% GA . 2% HQ, 10% GA			40	N/K	12	2% KA, 2% HQ, 10% GA P r 60% . 47.5% P r 2% HQ, 10% GA
F r r a C a r G r a . [84], [85] V r a -R a . [86]		R, O	4% HQ, 0.05% RA, 0.01% FA (TC) . 4% HQ + SPF 30			120	M/S	8	>75% P r , 73% TC/4 -R .9560414 c 83.6041-1.198801

T 3 Ac^u c f ra^p u c a r a a [86]

T 4ⁱ i

Tra a c ac c

V a C

V a E

S b a rac

T^p ca q^u r

L c r c rac

G a^ψ f

B a^b r r rac

P^p r^u b r rac

A^r Bⁱ

I c f ac

N ac a c

N c c ac c

4-N Bⁱ r r c r^u r

M a^ψ f

light (IPL), cryosurgery, dermabrasion, and micro-dermabrasion. They are not considered as first-line therapy as they can cause PIH and are often ineffective [54–56]. Chemical peels, lasers, and IPL are described here as their use in melasma is better studied in the Indian setting to some extent.

Chemical Peels

Chemical peels used for treating melasma are described in Table 4. There is a medium-quality evidence to suggest that undergoing serial chemical peels with GA, salicylic acid, or trichloroacetic acid (TCA) is moderately effective in the treatment of melasma (Table 5) [22, 53, 57, 58]. GA peels are used as an adjunctive therapy in refractory cases of epidermal melasma as they enhance the efficacy of topical regimen [59]. There are

many studies comparing GA and other peels for melasma in dark-skinned patients (Table 6) [53].

Chemical peels cause controlled exfoliation, followed by regeneration of epidermis and dermis [60]. Superficial and medium-depth peels have been employed with variable success in the treatment of melasma. After a superficial peel, epidermal regeneration can be expected within 3–5 days and desquamation is usually well accepted [61]. Because of the risk of prolonged dyschromia, medium-depth and deep peels should be avoided in patients with dark skin. In the authors' opinion, maintenance sessions or repeat sessions may be necessary. Patients should be well informed.

Lasers and Light Therapy

Lasers have a limited role in the treatment of melasma. Though successful use of Q-switched (QS) lasers [50], fractional lasers [57, 62–66], IPL [56], and combination lasers [67–69] has been reported, response to treatment is unpredictable and pigmentation frequently recurs. In addition, PIH is common in Indian patients. Hence, a maintenance schedule has to be initiated and continued. For these reasons, lasers are not routinely recommended as the treatment of choice for melasma in Indian patients. It may be used in selected resistant cases, at the discretion of the treating physician, after proper counseling. A test patch may be performed prior to treating the lesion. QS laser and low-fluence mode laser toning alone or in combination with either fractional CO₂ laser or IPL have shown reasonable success in treating melasma of Asian patients (Table 7). A number of studies have shown successful treatment of melasma with fractional lasers [non-ablative Er (Erbium): Glass 1,540/1,550 nm lasers], and they are recommended in all skin types [57, 62–65].

T	4	Tra	c a c r	r q	ca P
C	T	T	T	C	T
GA	30–70%	SUP r ca P	A r a P	, r a GA	–
		P ar a P c 5 5	r 2 3 W	. T P	
		u r a c u	a r r 1% b car b a		
		u	[53, 56, 58–60, 73, 87, 88]		
LA	92%	SUP r ca P	. 2 c a	LA a P H 3.5 ar	Sa a c c c , ac
		a P c r 10	r 3 W	[87–89]	
SA	SUP r ca P	. 20–30% P u	HQ a 2- W	r a T c c car r	c q r a ,
				u P r ca P	b u c W c a [62]. A W
				c r a	SA W a acc a a q a , r
				P r c ac c, r a c r a c P r c , a r	
				ar c q a	ac a c r a r r a c
				c . I	b c r a c r P
				q u a	c a c a r r a r c a SA
				P a	W a a [54, 55, 59, 60, 89, 90]
TCA	10–30%	SUP r ca P	. Or 35–50%	c u -c P r	Ma c a car r a c PIH car , b u c
		P . P	r 2 r , >13 r . P	W a r c	W c a [53, 58–60, 90]
		u P			
Tr	r 5%	SUP r ca P	. Tr	a 1% r r	–
		a P c r 4 r	c a W	, r 12 W	[91, 92]
MA	SUP r ca P	. MA a 30–50%	a P c W	r A a ab	a a r ac r ba a r 10%
		b W	a c u c a a ac	W a a 2%	
				a r c b a	W a C a c
				E . L	r r a a c r c c W a r [53, 93]
R c	A P c c a W	B u ca b r P a c W c a W	N u r a a	r q u r c . Sa a c c c	r car
ac c	r acc r a c c . 5–6 P	ar r q u r c	. N r a	, B r	, r car [53, 58, 94]
	a q r				
GA c c ac c, HQ r c q u , MA a c c ac c, LA Lac c ac c, PIH P a a r r P P a , SA a c c ac c, TCA r q r ac c ac c					

ALGORITHM FOR INDIAN MELASMA PATIENTS

India is a country of multiple ethnicity and origin, covering a wide range of skin types. Melasma is found more frequently in Fitzpatrick type III–VI skin [48]. Based on Pigmentary Disorders Academy (PDA) recommendations

for melasma treatment for dark-skin types and a thorough review and grading of the literature on evidence-based studies (listed in Table 8) [48, 61] an algorithm is proposed in Fig. 1 to treat Indian melasma patients. All treatments included and their hierarchy in the algorithm is based on the grading of the associated clinical studies according to the US preventive services

T	5 L	ca	ca	ca	ca	ca	ca	ca	ca
C	P	L	S	R					
S ^u P ^r ca P ^r	R ^r c ac c ^r	E ^r P ^r P ^r	C	[53]					
	Lac c ac c ^r	U c ^r P ^r c ^r a	B	[87, 88]					
	G c c ac c ^r	N ^r a c ^r c ^r	A	[55, 58, 59, 72, 89, 90, 94–96]					
	Sa c c ac c ^r	U c ^r P ^r c ^r a	B	[54]					
	J ^r u	U c ^r P ^r c ^r a	B	[97]					
S ^u P ^r ca- c ^u P ^r	P ^u c ac c ^r	E ^r P ^r P ^r	C	[98]					
M c ^u -c ^r P ^r P ^r	Tr ^r P ^r ac c ^r U c ^r P ^r c ^r a	B	[96, 97]						

A: T^r P^r c^r c^r c^r u P^r P^r u P^r P^r c^r c^r, B: T^r P^r a^r c^r c^r u P^r P^r u P^r P^r c^r c^r,
C: T^r P^r P^r c^r c^r u P^r P^r u P^r P^r c^r c^r

T	6 C	Para	u c ^r P ^r P ^r ca P ^r	ca P ^r	ca P ^r	ca P ^r	ca P ^r	ca P ^r	ca P ^r
R	S	T	P	T	R				
Ka a a. R, O	55–75% GA	100	E ^r 15 ca	R ^r a ^r TCA a c ^r P ^r a ^r c ^r	[89]				
K ^u a. [90]	O, SF	1% P ^r	10	B ^r 12 P ^r	D c ^r a c ^r MASI, c ^r c ^r				
S ^r a ^u a. [87, 88]	O, SF	92% P ^u ac c ac c ^r	30	E ^r 3 P ^r P ^r a ^u 5	S a ca ca P ^r b P ^r c ^r				
Sa ^u a. [97]	O, SF	M c ^r c ^r J ^r u + 15% TCA	20	E ^r 10 ca P ^r 8	B ^r P ^r c b a P ^r				
K ^u a ^r a c ^r T ^r a ^r P ^r a [94]	R, O	20–35% GA 10–20% TCA	40	E ^r 15 ca 12 P ^r	Ab ^u 75% P ^r b P ^r c ^r				
S ^r b ^r a c ^r S ^r b ^r [96]	O, SF	70% GA a C	14	6	B ^r P ^r u a C c ^r				

GA c c ac c^r P^r, MASI a a a^r a^r c^r, O P^r, R a c^r c^r, SF P^r ac, TCA P^r P^r ac c ac c^r P^r

T	7	Tra	c a c r	r a r f r a ^p				
T		M			T	f	f	C
QS Nd:YAG La r	1,064	(			10	,	c	E c ^p
u c c a r		a						c r f a,
)		a c a c						f f . A ar
		r a c . A						p (6) f u c b
		p f ac ^u c c . S ^u b-						p c Γ 2 p a f f
		c u ar c						a r a ^p . T f
		p f f r c c ^u r						r a r c c c
		f f u c c .						a p . Pr f f
		D r a f f u c						r p -c b a c r a
		ca a						p r r a r f r a ^p
								r c c c [91, 96]. PIH
								a c b f c c ^p a
								f a b r p r c [91, 99]
C b a QS	La r	, ^u	a ar ^p	10	,	r	2 3 ^p	Pr r a ca .
Nd:YAG 1,064, f f		f f r f u c						C c a a c ^p -
f r ac a CO ₂		u p p a a						f r a ^p p ca r a
a r		r q ^u r a						a a r .
								Ma a c a b
								c c ca r a ^p
IPL		Sa a a r						E c r a r r ac r
								a a A a p a
								[56, 67]
C b a IPL	Sa a a r			1	IPL r c ar			Ra ^p c r u c- ^p
f f QS Nd:YAG					p c r a a a f c			a a f f p b -
a r 1,064 (f					b 4 5 QS			r b [67 69]. T f
u c c a r					Nd:YAG a r a 2- ^p			r a r c c c
)					r a			a p
IPL								
QS Q								

task force levels of evidence for grading clinical trials [70].

The key considerations in developing the algorithm are: the Fitzpatrick skin types susceptible to melasma in the Indian population, the severity of melasma, the sensitivity of patients to active ingredients of medication, the patient treatment history, any existing skin condition (besides melasma), possible therapy-related AEs such as PIH, exogenous ochronosis, blotchy

depigmentation, irritation, erythema; and the probability of pigment recurrence after stopping treatment, i.e., prognosis indicators. The darker skin types tend to be highly sensitive to treatment agents, for example, ochronosis on HQ treatment, sensitization to KA, and skin irritation reactions to peeling agents like GA. In addition, darker skin types are more prone to PIH post-treatment and have a greater chance of relapse [71]. With these considerations in mind, darker skin types, which are commoner among

T 8 L a c q^u a c c r a a f a^p

T	L _v v ^l	Q _v ψ _v v ^l	R
T ^p ca			
2% HQ	II	C	[83]
4% HQ	I	B	[27, 43, 78, 85]
0.1% r (RA)	I	B	[20, 100]
0.05% RA	I	C	[92]
0.05% I r	II	C	[92, 101]
4% <i>N</i> -ac -4- <i>S</i> -C a P _f	III	C	[101, 102]
5% HQ + 0.1 0.4% RA + 7% LA/10% AC	III	C	[102, 103]
3% HQ + 0.1% RA	III	C	[102, 103]
2% HQ + 0.05% RA + 0.01% ^u c ac c	I	A	[38, 39, 41, 74, 85, 103, 104]
2% HQ + 0.05% RA + 0.01% c a f a (c c KF)	III	C	[55, 86]
2% HQ + 0.05% RA + 0.01% c a f a (c c KF + 30 40% GA ^p)	III	B	[55, 86]
5% HQ + 0.1% RA, a c 1% f c c r	III	C	[32, 33]
4% HQ + 5% GA	II	B	[104, 105]
4% KA + 5% GA	II	B	[98, 99]
2% KA + 2% HQ + 10% GA	II	C	[83, 84]
2% HQ + 10% GA	II	C	[83, 84]
4% HQ + 10% GA	I	B	[105, 106]
20% A a c ac c	I	B	[42, 43]
20% A a c ac c + 0.05% RA	III	C	[106, 107]
V a C P _f r	II	C	[107, 108]
Ac a ^p a	II	B	[99, 100]
Q _f ca ^p			
10 50% GA	II /III	C	[53, 59]
10% + 2% HQ + 20 70% GA	II	C	[56, 58]
20 30% GA + 1071100 1071100 56TD[(8339.6(A)-642705%)-339.6(RA)-12980.9003906(III)-5135.4(C)-6103.60009(I)],			

Table 8 c u c

T	L v v ¹	Q _u -ψ v ¹	R
50% GA + 10% KA La r k ra ^p (+ q ca ^p + p ca k ra ^p) Q- ^ψ q c ^u b p ^u c CO ₂ + Q- ^ψ q c a a c ^r Q- ^ψ q c a a c ^r a r Q- ^ψ q c a a c ^r a r + 15-25% TCA ^p + J r' u	III IV IV IV III	C C C C C	[109, 112] [110, 113] [67, 68] [69, 70] [111, 114]
Er ^b : YAG D r abra	III II	D E	[112, 115] [76, 113]
I acc rda c ^ψ k k US ^p r c a r c c c r rad c ca ra . R ^p c ^u c ^ψ k p r [48] A: k r c c c u p p r k u k p r c ^u r, B: k r a r c c u p p r k u k p r c ^u r, C: k r p r c c u p p r k u k p r c ^u r, D: k r a r c c u p p r k r c k u k p r c ^u r, E: k r c c c u p p r k r c k u k p r c ^u r AC			

Indian people, are recommended longer treatment periods, with lower concentrations of those treatment agents that can induce PIH or irritation, followed by longer maintenance periods to prevent recurrence of pigmentation and melasma. We have found that treatments like laser therapy, although described for all skin types in literature, are in fact not suitable for Indian skin as PIH and relapse occurs frequently once treatment is stopped in spite of maintenance therapy. However, light therapy does have some value in Indian patients.

In the opinion of the authors, broad-spectrum sunscreen use during and post-treatment is mandatory, along with protective clothing and hats. Experts recommend the fixed triple combination as the first-line therapy for all melasma types and degrees of severity, and dual combinations or single agents should be considered only when it is unavailable or

patients have sensitivity to the ingredients [48]. In such cases, we suggest dual-combination therapy (e.g., HQ + GA) or monotherapy (e.g., 4% HQ, 0.1% retinoic acid, or 20% AA). For moderate or severe melasma which does not respond to first-line treatment, options for second-line therapy include peels either alone or in combination with topical therapy.

In the opinion of the authors, some patients will require therapy to maintain remission status and a combination of topical therapies should be considered. A minimum of four sessions of a given peel are suggested before changing to an alternative therapy. It is to be noted that for Indian skin types, especially Fitzpatrick types IV–VI, the concentration of certain agents should be reduced; 5% HQ should be avoided for Indian patients as PIH and irritation are more likely at higher




Fig. 1 A $\times 10$ magnification of a skin biopsy specimen from a patient with melasma. The histological slide shows a cross-section of skin with increased melanin in the epidermis and dermis. The clinical photo shows dark, irregular pigmentation on the forehead and cheeks.

concentrations. Because of the risk of prolonged hyperpigmentation, medium-depth peels like $\geq 35\%$ TCA should be conducted with caution in patients with dark-skin types; deep peels should not be used in these patients. Furthermore, chemical peels are generally used to treat only the epidermal and mixed forms of melasma, as an attempt to treat with deep peels often leads to unwanted complications like hypertrophic scarring and permanent depigmentation. There are ample studies on the effect of GA peels on ethnic skin [55] (Table 9) and hence make for a recommended choice of treatment. In most of these studies, there was a moderate improvement achieved in almost one-half of the patients [58, 59, 72, 73].

Lasers should rarely be used in the treatment of melasma and, if applied, it is important to consider skin type. The mode-toning laser treatment is generally recommended in all skin types and its major advantage is that it destroys melanin without cell damage. Both QS low fluence and IPL are effective for recalcitrant melasma, when patients are non-responsive to other treatments, or when patients are intolerant of other treatment medications. The combination of IPL with QS laser gives a rapid resolution of mixed-type melasma with possible long-term benefits [67–69]. Fractional non-ablative lasers can be considered only if all other modalities fail. Ablative lasers should not be considered.

T 9 S^u c a a f a^p f c [53]

R	N	P	T	R
L a c f a	10 A a ()	20 70% –		N a ca ca
[58]		GA		
G r [54]	6 (car r a c a f c	20 30% –		M c r a p r 66% p a
	r u p USA)	SA		
J a f a r a .	25 I c a	50% GA S ^u c r SPF 15,		46.7% p c r a , 27.8% c , 0%
[59]		10% GA		c r a
G r a c	15 I c a	S r a GA –		G c a r p r
R c c ^u [73]		p		
S a r ^u a .	20 I r a	92% LA –		S ca p r a 12
[87]				p a f c p c u c
G c a c	20 I c a	S r a GA T r p c b a		>50% p r f a p a
S a f a [72]		p a c u c r		

GA c c a c c, LA a c c a c c, SA a c c a c c

Patient consent and a photograph of the condition must be obtained prior to therapy. As important as actual therapy is proper patient counseling, and an understanding of the psychosocial impact on the patient and of his/her condition leading to low quality of life [74, 75]. This is likely to lead to a better patient compliance and an adherence to the regimen. It has been reported in Indian patients that overuse of certain treatment modalities such as mometasone-based therapies leads to undesirable fallouts [47, 49].

The recommendation is to use an alternative triple-combination cream, better suited for Indian patients, i.e., 2% HQ, 0.05% tretinoin, and 0.01% of the mild steroid fluocinolone acetonide cream. Another option is to replace the mometasone-based triple combination after 4–8 weeks, with a fluocinolone or a hydrocortisone-based one for a further 3- to 6-month period. As discussed earlier, this cream is safe and effective based on extensive clinical studies. The gradual withdrawal of the

T	10 Bac p r a c r
F	f f f v f v f f f f
f p III VI: c a r f a r a c / r c a r	
G c a c a a p r c p [3, 6, 8 10]	
L - r a a p ≥2 a r r a	
H r p r c u r a r	
T r a c b ≥2 p f c a	
L - r - r a f r c [46, 47]	
O f r [76, 114]	
M c- p a a [115]	

therapeutic agent is also a consideration and is achieved by decreasing doses of the active agent. The proposed algorithm for Indian patients is presented in Fig. 1.

BAD PROGNOSIS INDICATORS

There are several conditions that determine the outcome of treatment in melasma since its

etiology is multifactorial. Several factors predict potential failure of treatment (listed in Table 10). Moin et al. [3] found a statistically significant relationship between melasma and ethnicity, and phototype and grade of parity. Exogenous ochronosis has also been reported to be being misdiagnosed as a melasma treatment failure [76]. However, these prognostic factors lack sufficient scientific evidence.

CONCLUSION

Here, the Indian dermatologists panel has graded the results of safety and efficacy of various melasma therapies from clinical trials conducted worldwide (Table 7) applying the quality and level

3. Moin A, Jabery Z, Fallah N. Prevalence and awareness of melasma during pregnancy. *Int J Dermatol*. 2006;45(3):285–8 (Epub 2006/03/15).
4. Kim NH, Lee CH, Lee AY. H19 RNA downregulation stimulated melanogenesis in melasma. *Pigment Cell Melanoma Res*. 2010;23(1):84–92 (Epub 2009/12/09).
5. Draelos ZD. Skin lightening preparations and the hydroquinone controversy. *Dermatol Ther*. 2007;20(5):308–13 (Epub 2007/11/30).
6. Goh CL, Dlova CN. A retrospective study on the clinical presentation and treatment outcome of melasma in a tertiary dermatological referral centre in Singapore. *Singapore Med J*. 1999;40(7):455–8 (Epub 1999/11/24).
7. Ortonne JP, Arellano I, Berneburg M, et al. A global survey of the role of ultraviolet radiation and hormonal influences in the development of melasma. *J Eur Acad Dermatol Venereol*. 2009;23(11):1254–62 (Epub 2009/06/03).
8. Sarkar R, Puri P, Jain RK, Singh A, Desai A. Melasma in men: a clinical, aetiological and histological study. *J Eur Acad Dermatol Venereol*. 2010;24(7):768–72 (Epub 2009/12/18).
9. Vazquez M, Maldonado H, Benmaman C, Sanchez JL. Melasma in men. A clinical and histologic study. *Int J Dermatol*. 1988;27(1):25–7 (Epub 1988/01/01).
10. Sivayathorn A. Melasma in orientals. *Clin Drug Invest*. 1995;10(Suppl. 2):34–40.
11. Sarvjot V, Sharma S, Mishra S, Singh A. Melasma: a clinicopathological study of 43 cases. *Indian J Pathol Microbiol*. 2009;52(3):357–9 (Epub 2009/08/15).
12. Grimes PE, Yamada N, Bhawan J. Light microscopic, immunohistochemical, and ultrastructural alterations in patients with melasma. *Am J Dermatopathol*. 2005;27(2):96–101 Epub 2005/03/31.
13. Kang WH, Yoon KH, Lee ES, et al. Melasma: histopathological characteristics in 56 Korean patients. *Br J Dermatol*. 2002;146(2):228–37 (Epub 2002/03/21).
14. Verallo-Rowell VM. Skin in the tropics: Sunscreen and hyperpigmentation. Philippines: Anvil

- Clin Dermatol. 2001;19(4):483–8 (Epub 2001/09/06).
29. Prignano F, Ortonne JP, Buggiani G, Lotti T. Therapeutical approaches in melasma. *Dermatol Clin*. 2007;25(3):337–42 (viii. Epub 2007/07/31).
 30. Bandyopadhyay D. Topical treatment of melasma. *Indian J Dermatol*. 2009;54(4):303–9 (Epub 2010/01/27).
 31. Lynde CB, Kraft JN, Lynde CW. Topical treatments for melasma and postinflammatory hyperpigmentation. *Skin Ther Lett*. 2006;11(9):1–6 (Epub 2006/11/01).
 32. Kligman AM, Willis I. A new formula for depigmenting human skin. *Arch Dermatol*. 1975;111(1):40–8 (Epub 1975/01/01).
 33. Kang WH, Chun SC, Lee S. Intermittent therapy for melasma in Asian patients with combined topical agents (retinoic acid, hydroquinone and hydrocortisone): clinical and histological studies. *J Dermatol*. 1998;25(9):587–96 (Epub 1998/11/03).
 34. Godse KV. Triple combination of hydroquinone, tretinoin and mometasone furoate with glycolic acid peels in melasma. *Indian J Dermatol*. 2009;54(1):92–3 (Epub 2010/01/06).
 35. Kulhalli P, Chevli T, Karnik R, Sheth M, Mulgaonkar N. Comparative potency of formulations of mometasone furoate in terms of inhibition of 'PIRHR' in the forearm skin of normal human subjects measured with laser doppler velocimetry. *Indian J Dermatol Venereol Leprol*. 2005;71(3):170–4 (Epub 2006/01/06).
 36. Chan R, Park KC, Lee MH, et al. A randomized controlled trial of the efficacy and safety of a fixed triple combination (fluocinolone acetonide 0.01%, hydroquinone 4%, tretinoin 0.05%) compared with hydroquinone 4% cream in Asian patients with moderate to severe melasma. *Br J Dermatol*. 2008;159(3):697–703 (Epub 2008/07/12).
 37. Jamkhedkar P, Shenai C, Shroff HJ, et al. Fluticasone propionate (0.05%) cream compared to betamethasone valerate (0.12%) cream in the treatment of steroid-responsive dermatoses: a multicentric study. *Indian J Dermatol Venereol Leprol*. 1996;62(5):289–94 (Epub 1996/09/01).
 38. Rajaratnam R, Halpern J, Salim A, Emmett C. Interventions for melasma. *Cochrane Database Syst Rev*. 2010(7):CD003583. (Epub 2010/07/09).
 39. Taylor SC, Torok H, Jones T, et al. Efficacy and safety of a new triple-combination agent for the treatment of facial melasma. *Cutis*. 2003;72(1):67–72 (Epub 2003/08/02).
 40. Torok H, Taylor S, Baumann L, et al. A large 12-month extension study of an 8-week trial to evaluate the safety and efficacy of triple combination (TC) cream in melasma patients previously treated with TC cream or one of its dyads. *J Drugs Dermatol*. 2005;4(5):592–7 (Epub 2005/09/20).
 41. Torok HM. A comprehensive review of the long-term and short-term treatment of melasma with a triple combination cream. *Am J Clin Dermatol*. 2006;7(4):223–30 (Epub 2006/08/12).
 42. Nazzaro-Porro M. Azelaic acid. *J Am Acad Dermatol*. 1987;17(6):1033–41 (Epub 1987/12/01).
 43. Balina LM, Graupe K. The treatment of melasma. 20% azelaic acid versus 4% hydroquinone cream. *Int J Dermatol*. 1991;30(12):893–5 (Epub 1991/12/01).
 44. Godse DK. Approach to melasma in India. Available from: http://www.academia.edu/3647500/approach_to_melasma_in_india. Accessed Sept 8, 2014.
 45. Marta Rendon MB, Ivonne Arellano, Mauro Picardo, et al. Treatment of melasma. Available from: <http://leenyx.co.za/backend/media/54201093909AM/Treatment%20of%20melasma.pdf>. Accessed Sept 8, 2014.
 46. Sehgal VN, Verma P, Srivastava G, Aggarwal AK, Verma S. Melasma: treatment strategy. *J Cosmet Laser Ther*. 2011;13(6):265–79 (Epub 2011/10/11).
 47. Majid I. Mometasone-based triple combination therapy in melasma: is it really safe? *Indian J Dermatol*. 2010;55(4):359–62 (Epub 2011/03/25).
 48. Rendon M, Berneburg M, Arellano I, Picardo M. Treatment of melasma. *J Am Acad Dermatol*. 2006;54(5 Suppl 2):S272–81 (Epub 2006/04/25).
 49. Godse KV, Zawar V. Mometasone menace in melasma. *Indian J Dermatol*. 2012;57(4):324–6 (Epub 2012/07/28).
 50. Wu S, Shi H, Wu H, et al. Treatment of melasma with oral administration of tranexamic acid. *Aesthetic Plast Surg*. 2012;36(4):964–70 (Epub 2012/05/04).
 51. Kanechorn Na Ayuthaya P, Niumphradit N, Manosroi A, Nakakes A. Topical 5% tranexamic acid for the treatment of melasma in Asians: a double-blind randomized controlled clinical trial. *J Cosmet Laser Ther*. 2012;14(3):150–4 (Epub 2012/04/18).

52. Lee JH, Park JG, Lim SH, et al. Localized intradermal microinjection of tranexamic acid for treatment of melasma in Asian patients: a preliminary clinical trial. *Dermatol Surg.* 2006;32(5):626–31 (Epub 2006/05/19).
53. Sarkar R, Bansal S, Garg VK. Chemical peels for melasma in dark-skinned patients. *J Cutan Aesthet Surg.* 2012;5(4):247–53 (Epub 2013/02/05).
54. Grimes PE. The safety and efficacy of salicylic acid chemical peels in darker racial-ethnic groups. *Dermatol Surg.* 1999;25(1):18–22 (Epub 1999/02/06).
55. Sarkar R, Kaur C, Bhalla M, Kanwar AJ. The combination of glycolic acid peels with a topical regimen in the treatment of melasma in dark-skinned patients: a comparative study. *Dermatol Surg.* 2002;28(9):828–32 (discussion 32. Epub 2002/09/25).
56. Wang CC, Hui CY, Sue YM, Wong WR, Hong HS. Intense pulsed light for the treatment of refractory melasma in Asian persons. *Dermatol Surg.* 2004;30(9):1196–200 (Epub 2004/09/10).
57. Naito SK. Fractional photothermolysis treatment for resistant melasma in Chinese females. *J Cosmet Laser Ther.* 2007;9(3):161–3 (Epub 2007/09/01).
58. Lim JT, Tham SN. Glycolic acid peels in the treatment of melasma among Asian women. *Dermatol Surg.* 1997;23(3):177–9 (Epub 1997/03/01).
59. Javaheri SM, Handa S, Kaur I, Kumar B. Safety and efficacy of glycolic acid facial peel in Indian women with melasma. *Int J Dermatol.* 2001;40(5):354–7 (Epub 2001/09/14).
60. Khunger N. Standard guidelines of care for chemical peels. *Indian J Dermatol Venereol Leprol.* 2008;74(Suppl):S5–12 (Epub 2008/08/21).
61. Rendon MI, Berson DS, Cohen JL, Roberts WE, Starker I, Wang B. Evidence and considerations in the application of chemical peels in skin disorders and aesthetic resurfacing. *J Clin Aesthet Dermatol.* 2010;3(7):32–43 (Epub 2010/08/21).
62. Goel A, Krupashankar DS, Aurangabadkar S, Nischal KC, Omprakash HM, Mysore V. Fractional lasers in dermatology—current status and recommendations. *Indian J Dermatol Venereol Leprol.* 2011;77(3):369–79 (Epub 2011/04/22).
63. Goldberg DJ, Berlin AL, Phelps R. Histologic and ultrastructural analysis of melasma after fractional resurfacing. *Lasers Surg Med.* 2008;40(2):134–8 (Epub 2008/02/29).
64. Rokhsar CK, Fitzpatrick RE. The treatment of melasma with fractional photothermolysis: a pilot study. *Dermatol Surg.* 2005;31(12):1645–50 (Epub 2005/12/13).
65. Tannous ZS, Astner S. Utilizing fractional resurfacing in the treatment of therapy-resistant melasma. *J Cosmet Laser Ther.* 2005;7(1):39–43 (Epub 2005/07/16).
66. Trelles MA, Velez M, Gold MH. The treatment of melasma with topical creams alone, CO2 fractional ablative resurfacing alone, or a combination of the two: a comparative study. *J Drugs Dermatol.* 2010;9(4):315–22 (Epub 2010/06/03).
67. Na SY, Cho S, Lee JH. Intense pulsed light and low-fluence Q-switched Nd:YAG laser treatment in melasma patients. *Ann Dermatol.* 2012;24(3):267–73 (Epub 2012/08/11).
68. Nouri K, Bowes L, Chartier T, Romagosa R, Spencer J. Combination treatment of melasma with pulsed CO2 laser followed by Q-switched alexandrite laser: a pilot study. *Dermatol Surg.* 1999;25(6):494–7 (Epub 1999/09/01).
69. Angsuwarangsee S, Polnikorn N. Combined ultrapulse CO2 laser and Q-switched alexandrite laser compared with Q-switched alexandrite laser alone for refractory melasma: split-face design. *Dermatol Surg.* 2003;29(1):59–64 (Epub 2003/01/22).
70. Rockville. Appendix A How the U.S. Preventive Services Task Force Grades Its Recommendations. The Guide to Clinical Preventive Services 2012: Recommendations of the U.S. Preventive Services Task Force. 2012. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK115122/>. Accessed Sept, 9 2014.
71. Lin JY, Chan HH. Pigmentary disorders in Asian skin: treatment with laser and intense pulsed light sources. *Skin Ther Lett.* 2006;11(8):8–11 (Epub 2006/10/07).
72. Godse KV, Sakhia J. Triple combination and glycolic acid peels in melasma in Indian patients. *J Cosmet Dermatol.* 2011;10(1):68–9 (Epub 2011/02/22).
73. Grover C, Reddu BS. The therapeutic value of glycolic acid peels in dermatology. *Indian J Dermatol Venereol Leprol.* 2003;69(2):148–50 (Epub 2007/07/24).
74. Cestari TF, Hexsel D, Viegas ML, et al. Validation of a melasma quality of life questionnaire for Brazilian Portuguese language: the MelasQoL-BP study and improvement of QoL of melasma patients after

- triple combination therapy. *Br J Dermatol*. 2006;156(Suppl 1):13–20 (Epub 2006/12/21).
75. Pichardo R, Vallejos Q, Feldman SR, et al. The prevalence of melasma and its association with quality of life in adult male Latino migrant workers. *Int J Dermatol*. 2009;48(1):22–6 (Epub 2009/01/08).
 76. Ribas J, Schettini AP, Cavalcante Mde S. Exogenous ochronosis hydroquinone induced: a report of four cases. *An Bras Dermatol*. 2010;85(5):699–703 (Epub 2010/12/15).
 77. Monteiro RC, Kishore BN, Bhat RM, Sukumar D, Martis J, Ganesh HK. A comparative study of the efficacy of 4% hydroquinone vs 0.75% Kojic acid cream in the treatment of facial melasma. *Indian J Dermatol*. 2013;58(2):157.
 78. Farshi S. Comparative study of therapeutic effects of 20% azelaic acid and hydroquinone 4% cream in the treatment of melasma. *J Cosmet Dermatol*. 2011;10(4):282–7 (Epub 2011/12/14).
 79. Espinal-Perez LE, Moncada B, Castanedo-Cazares JP. A double-blind randomized trial of 5% ascorbic acid vs. 4% hydroquinone in melasma. *Int J Dermatol*. 2004;43(8):604–7 (Epub 2004/08/12).
 80. Nanda S, Grover C, Reddy BS. Efficacy of hydroquinone (2%) versus tretinoin (0.025%) as adjunct topical agents for chemical peeling in patients of melasma. *Dermatol Surg*. 2004;30(3):385–8 (discussion 9. Epub 2004/03/11).
 81. Piquero Martin J, de Rothe Arocha J, Beniamini Loker D. Double-blind clinical study of the treatment of melasma with azelaic acid versus hydroquinone. *Med Cutan Ibero Lat Am*. 1988;16(6):511–4 (Epub 1988/01/01).
 82. Verallo-Rowell VM, Verallo V, Graupe K, Lopez-Villafuerte L, Garcia-Lopez M. Double-blind comparison of azelaic acid and hydroquinone in the treatment of melasma. *Acta Derm Venereol Suppl (Stockh)*. 1989;143:58–61 (Epub 1989/01/01).
 83. Lim JT. Treatment of melasma using kojic acid in a gel containing hydroquinone and glycolic acid. *Dermatol Surg*. 1999;25(4):282–4 (Epub 1999/07/27).
 84. Ferreira Cestari T, Hassun K, Sittart A, de Lourdes Viegas M. A comparison of triple combination cream and hydroquinone 4% cream for the treatment of moderate to severe facial melasma. *J Cosmet Dermatol*. 2007;6(1):36–9 (Epub 2007/03/14).
 85. Grimes P, Kelly AP, Torok H, Willis I. Community-based trial of a triple-combination agent for the treatment of facial melasma. *Cutis*. 2006;77(3):177–84 (Epub 2006/04/14).
 86. Rothe de Arocha J. Nuevas opciones en al tratamiento del melasma. *Dermatologia Venezolana*. 2003;41:11–4. Available from <http://revista.svderma.org/index.php/ojs/article/view/285/285>. Accessed Sept 9, 2014.
 87. Sharquie KE, Al-Tikreety MM, Al-Mashhadani SA. Lactic acid as a new therapeutic peeling agent in melasma. *Dermatol Surg*. 2005;31(2):149–54 (discussion 54. Epub 2005/03/15).
 88. Sharquie KE, Al-Tikreety MM, Al-Mashhadani SA. Lactic acid chemical peels as a new therapeutic modality in melasma in comparison to Jessner's solution chemical peels. *Dermatol Surg*. 2006;32(12):1429–36 (Epub 2007/01/04).
 89. Kalla G, Garg A, Kachhawa D. Chemical peeling—glycolic acid versus trichloroacetic acid in melasma. *Indian J Dermatol Venereol Leprol*. 2001;67(2):82–4 (Epub 2007/08/01).
 90. Khunger N, Sarkar R, Jain RK. Tretinoin peels versus glycolic acid peels in the treatment of Melasma in dark-skinned patients. *Dermatol Surg*. 2004;30(5):756–60 (discussion 60. Epub 2004/04/22).
 91. Griffiths CE, Finkel LJ, Ditre CM, Hamilton TA, Ellis CN, Voorhees JJ. Topical tretinoin (retinoic acid) improves melasma. A vehicle-controlled, clinical trial. *Br J Dermatol*. 1993;129(4):415–21 (Epub 1993/10/01).
 92. Jimbow K. N-acetyl-4-S-cysteaminylphenol as a new type of depigmenting agent for the melanoderma of patients with melasma. *Arch Dermatol*. 1991;127(10):1528–34 (Epub 1991/10/01).
 93. Taylor MB. Summary of mandelic acid for the improvement of skin conditions. *Cosmet Dermatol*. 1999;12:26–8.
 94. Kumari R, Thappa DM. Comparative study of trichloroacetic acid versus glycolic acid chemical peels in the treatment of melasma. *Indian J Dermatol Venereol Leprol*. 2010;76(4):447 (Epub 2010/07/27).
 95. Grimes PE. Melasma. Etiologic and therapeutic considerations. *Arch Dermatol*. 1995;131(12):1453–7 (Epub 1995/12/01).
 96. Sobhi RM, Sobhi AM. A single-blinded comparative study between the use of glycolic acid 70% peel and the use of topical nanosome vitamin C iontophoresis in the treatment of melasma. *J Cosmet Dermatol*. 2012;11(1):65–71 (Epub 2012/03/01).

97. Safoury OS, Zaki NM, El Nabarawy EA, Farag EA. A study comparing chemical peeling using modified Jessner's solution and 15% trichloroacetic Acid versus 15% trichloroacetic acid in the treatment of melasma. *Indian J Dermatol.* 2009;54(1):41–5 (Epub 2010/01/06).
98. Berardesca E, Cameli N, Primavera G, Carrera M. Clinical and instrumental evaluation of skin